arXiv:1207.6396v1 [hep-ph] 26 Jul 2012

The effect of plasmon mass on spin light of neutrino in dense matter

A.Grigoriev[a], A.Lokhov[b], A.Studenikin[b,c] [1], A.Ternov[d]

[a] *Skobeltsyn Institute of Nuclear Physics, Moscow State University, 119991 Moscow, Russia*
[b] *Department of Theoretical Physics, Moscow State University, 119991 Moscow, Russia*
[c] *Joint Institute for Nuclear Research, 141980 Dubna, Russia*
[d] *Department of Theoretical Physics, Moscow Institute for Physics and Technology, 141700 Dolgoprudny, Russia*

Abstract

We develop the theory of spin light of neutrino in matter ($SL\nu$) and include the effect of plasma influence on the emitted photon. We use the special technique based on exact solutions of particles wave equations in matter to perform all the relevant calculations, and track how the plasmon mass enters the process characteristics including the neutrino energy spectrum, $SL\nu$ rate and power. The new feature it induces is the existence of the process threshold for which we have found the exact expression and the dependence of the rate and power on this threshold condition. The $SL\nu$ spatial distribution accounting for the above effects has been also obtained. These results might be of interest in connection with the recently reported hints of ultra-high energy neutrinos $E = 1 \div 10$ PeV observed by IceCube.

1. Introduction

Neutrino physics in matter and external electromagnetic fields is a rather longstanding research field nevertheless still having advances and providing some interesting predictions for various phenomena. A broad spectrum of issues here are connected with possible electromagnetic properties of neutrino (for more details refer to [1]). The recent studies of neutrino electromagnetic properties revealed a new mechanism of electromagnetic radiation by a neutrino propagating in dense matter that has been proposed in [2]. This type of electromagnetic radiation was called the spin light of neutrino in matter ($SL\nu$). In a quasi-classical treatment this radiation originates due to neutrino electromagnetic moment precession in dense background matter. The quantum theory of this phenomena has been developed in [3, 4].

A new convenient and elegant way of description of neutrino interaction processes in matter has been proposed and developed in a series of papers [3, 5] (see also [4]). The elaborated method is based on the use of the exact solutions of the modified Dirac equation

---

[1] studenik@srd.sinp.msu.ru

for a neutrino (the method of exact solutions) in matter. The method was used to describe the phenomenon of the spin light of neutrino in matter within quantum approach. Note that the most recent development of the method in application to studies of neutrino motion in different extended environments can be found in [6].

The $SL\nu$ is a process of photon emission in neutrino transition between different helicity states in matter. As it has been shown [3], in the relativistic regime $SL\nu$ mechanism could provide up to one half of the initial neutrino energy transition to the emitted radiation. It was also shown that the $SL\nu$ provides the spin polarization effect of neutrino beam moving in matter (similarly to the well-known effect of the electron spin self-polarization in synchrotron radiation [7]).

In this letter we make a reasonable step towards the completeness of the physical picture and consider the question of the plasmon mass influence on $SL\nu$. The estimations already performed in [8] indicated that the plasmon has a considerable mass that can affect the physics of the process. To see how the plasmon mass enters the $SL\nu$ quantities we appeal to the method of exact solutions and carry out all the computations relevant to $SL\nu$ in an exact way and track how the plasmon mass enters the process characteristics including the neutrino energy spectrum, $SL\nu$ rate and power with particular focus on the spatial distribution.

2. Kinematics of the process

First we briefly consider the external conditions in which the $SL\nu$ process is possible. Suppose the matter density is close to that of nuclear matter. In astrophysics it could be found, for example, in neutron stars with density span up to $10^{38-41} cm^{-3}$ [9]). As it has been shown in ([3]) in the case when neutrons are the main component in the background matter, in fact, an antineutrino (not neutrino) produce the $SL\nu$. Nevertheless we shall call it neutrino for convenience (the relevant formulae remain the same).

In turn, the plasmon effects emerge only due to the electron matter component which is considerably less dense in typical matter of a neutron star. For definiteness, in what follows, in respect of densities we assume the condition

$$n_e \simeq 0.1 n_n \, . \tag{1}$$

Due to the smallness of the charged fraction we neglect its influence on neutrinos and account below only for the neutron matter component in any quantities associated with neutrino.

For the description of neutrino propagation in matter we use the exact solutions of the modified Dirac equation for the neutrino field in matter [3]

$$\left\{ i\gamma_\mu \partial^\mu - \frac{1}{2}\gamma_\mu (1+\gamma^5) f^\mu - m_\nu \right\} \Psi(x) = 0, \tag{2}$$

where in the case of the particle motion through the non-moving and unpolarized neutron matter $f^\mu = -G_F/\sqrt{2}\,(n_n, \mathbf{0})$. With that equation (2) has plane-wave solution determined by the 4-momentum $p^{\mu} = (E, \mathbf{p})$ and quantum numbers of the helicity $s = \pm 1$ and the sign of energy $\varepsilon = \pm 1$. The corresponding neutrino energy is given by

$$E = \sqrt{(p + s\tilde{n})^2 + m_{\nu}^2} + \tilde{n}, \quad \tilde{n} = \frac{1}{2\sqrt{2}} G_F n_n. \tag{3}$$

The neutrino-photon coupling is described by the usual dipole electromagnetic vertex $\mathbf{\Gamma} = i\omega\big\{\big[\mathbf{\Sigma} \times \boldsymbol{\varkappa}\big] + i\gamma^5\mathbf{\Sigma}\big\}$, for details see [3, 5]. From the energy-momentum conservation one obtains

$$E = E' + \omega; \quad \mathbf{p} = \mathbf{p}' + \mathbf{k}, \tag{4}$$

where the primed values correspond to the outgoing neutrino. One has to account for the plasma dispersion law in the form of

$$\omega = \sqrt{k^2 + m_{\gamma}^2}, \quad m_{\gamma} = \sqrt{2\alpha}(3\sqrt{\pi} n_e)^{1/3}. \tag{5}$$

We consider the relativistic energy limit and $s = -1, s' = 1$ that is the most relevant case for applications. From the energy-momentum conservation (4) we obtained that

$$\omega\sqrt{(p + \widetilde{n})^2 + m_{\nu}^2} = \widetilde{n}(p' + p) + pk\cos\theta + \frac{m_{\gamma}^2}{2}, \tag{6}$$

where θ is the angle between the directions of the initial neutrino propagation and that of the radiated photon. Note that the structure of eq.(6) is very much similar to the corresponding equation for the case of the $SL\nu$ in matter with different masses of neutrinos in the initial and final state [10].

To obtain the threshold condition for the $SL\nu$ process one should consider the system (4) relative to the vector p'. The corresponding exact equation for p' is quadratic and thus can be resolved exactly [11]. The resulting threshold condition can be written in the form

$$\frac{m_{\gamma}^2 + {}^1\!/_2 m_{\gamma} m_{\nu}}{4\widetilde{n}p} < 1. \tag{7}$$

Note that the condition nontrivially depends on matter density through $\widetilde{n}$ and m_{γ} given by (3) and (5) respectively.

To meet the requirements on the parameters imposed by the threshold condition (7) it is worth to begin with considering the neutrino mass as its value is very restricted experimentally. To date, the most conservative upper limit for the neutrino mass which agrees with existing experiments is set at the order of 1 eV (for the most recent result from the direct search for neutrino mass refer to [12], for a major review of the field see for instance [13]). On the other hand, as it was shown in previous investigations [3, 5] that SL_{ν} is more efficient for high the matter density. Therefore we consider the highest densities of matter discussed in the literature, that are of the order 10^{38-41} cm^{-3} for nuclear matter [9, 14]. Indicated values correspond to the range of the "density parameter" $\widetilde{n} \simeq 1 - 10^3$ eV. Taking also into account the condition (1), from (5) we estimate the value of plasmon mass as $m_{\gamma} \simeq 10^{7-8}$ eV.

With the above scales of the parameters the term that contains neutrino mass in the numerator in Eq.(7) is not important and the threshold condition (7) is reduced to

$$m_\gamma^2/4\widetilde{n}p < 1. \tag{8}$$

That leads, together with the scales of the parameters mentioned above, to the allowable range for the neutrino momentum $p \gtrsim 10^{12}$ eV. Thus, even in the case the threshold condition is accounted for, the room for $SL\nu$ is opened for high energy neutrinos (see also [15]). The lower the density, the higher values for the neutrino momentum are needed for the $SL\nu$ process to be opened. The last relation corresponds to the case of highest matter density among considered, $n \simeq 10^{41}$ cm^{-3} (the dependance of the plasmon mass on the matter density is rather weak).

Note that in spite of the considered ultra-high neutrino energies the need for accounting for non-local contributions in the neutrino-matter interactions (emphasized earlier, for instance, in [8]) is not justified (see also [15]).

3. The rate and power of the $SL\nu$ process

Performing calculations similar to those described in [3] with including additionally the effects of plasma discussed above we arrive at the following expression for the $SL\nu$ radiation power

$$\Gamma = \frac{\mu^2}{2\pi} \int \frac{k^2}{\omega} S(k)\, \delta(E - E' - \omega)\, dk d\Omega, \tag{9}$$

$$S(k) = (k^2 + \omega^2) \left(1 - \frac{k - p\cos\theta}{p'} \cos\theta\right) - 2\omega k \left(\cos\theta - \frac{k - p\cos\theta}{p'}\right). \tag{10}$$

Here μ is the neutrino magnetic moment and θ is the angle between the direction of the initial neutrino and the plasmon momentum, $\mathbf{p}$ and $\mathbf{k}$ respectively.

The exact evaluation of the integral in (9) accounting for the composite delta function in the presented form is a rather involved procedure (also has been performed by the authors) and resulting expression is cumbersome to be presented here. Therefore it is reasonable to consider several ranges of parameters and obtain much simpler though less general expressions, that cover the whole parameters range of interest for astrophysical applications. Accounting for the threshold condition (8) it is convenient to consider the following three ranges of parameters and the corresponding three cases.

First we single out the "near-threshold" case,

$$m_\gamma^2/4\widetilde{n}p \lesssim 1. \tag{11}$$

Introducing notation $a = m_\gamma^2/4\widetilde{n}p$, the expression for the process transition rate, that follows from (11), can be written as

$$\Gamma = 4\mu^2\, \widetilde{n}^2 p \big((1-a)(1+7a) + 4a(1+a)\ln a\big), \tag{12}$$

in which one can recognize the estimation from [8]. For the corresponding expression for the total radiation power ($I = \int \omega d\Gamma$) our calculations yield

$$I = \frac{4}{3}\mu^2\widetilde{n}^2p^2(1-a)(1-5a-8a^2) - m_\gamma^4 \ln a. \tag{13}$$

The next case we consider is the "far-from-threshold" regime, realized when

$$m_\gamma^2/4\widetilde{n}p \ll 1, \quad \text{or} \quad a \to 0. \tag{14}$$

Since the value for $\widetilde{n}$ is restricted by $\widetilde{n} \simeq 1 - 10^3$ eV, the validity of condition (14) can be provided by the increase of the neutrino momentum alone, i.e. $p > 10^{12}$ eV. Thus this particular case, settled by the condition (14) can be referred to as the case of "ultra-relativistic" neutrinos. In this case the threshold effect is negligible so that the corresponding result of calculations without account for plasmon mass should be relevant. Indeed the expressions (12) and (13) under the condition (14) are transformed respectively to

$$\Gamma = 4\mu^2\widetilde{n}^2p, \tag{15}$$

$$I = \frac{4}{3}\mu^2\widetilde{n}^2p^2, \tag{16}$$

which are just results for the $SL\nu$ rate and power at high neutrino momentum p without inclusion of plasmon mass m_γ (see [3, 5]).

Now let us consider the case of the $SL\nu$ at the threshold, i.e. very close to it. In the case the following condition is realized

$$1 - m_\gamma^2/4\widetilde{n}p = 1 - a \ll 1, \quad \text{or} \quad a \to 1, \tag{17}$$

and the transition rate and radiation power are respectively given by

$$\Gamma = 4\mu^2\widetilde{n}^2(1-a)\left((1-a)p + 2\widetilde{n}\right), \tag{18}$$

$$I = 4\mu^2\widetilde{n}^2p(1-a)\left((1-a)p + 2\widetilde{n}\right). \tag{19}$$

The obtained expressions enable us to follow these two characteristics dependance on the neutrino energy as the parameter a reaches the threshold, $a \to 1$. As is expected the rate and power given by (18), (19) go to zero at the threshold.

The performed above analysis of the $SL\nu$ rate and power enable us to obtain a fine description for the angular distribution of the radiation. As follows from (9), (10) the radiation is confined within the narrow solid angle defined by the relation

$$\sin\theta \approx \theta \leq \frac{4\widetilde{n}(p-\widetilde{n}) - m_\gamma^2}{2pm_\gamma}. \tag{20}$$

The angular distribution of the $SL\nu$ radiation is presented on Fig.1 and Fig.2.

Far from the threshold (at high values of the neutrino momentum p) the angular distribution of the $SL\nu$ exhibit the characteristic that are typical for the radiation produced



Figure 1: 3D representation of the radiation power distribution.



Figure 2: The two-dimensional cut along the symmetry axis. Relative units are used.

by a relativistic particle. As is clearly seen on Fig.1 and Fig.2, the $SL\nu$ radiation is focused in the direction of the initial neutrino propagation in a very narrow solid angle $\theta \lesssim 2\widetilde{n}/m_\gamma$. The radiation distribution has a cup-like shape at the directions closest to the initial neutrino momentum. A remarkable feature that one can observe on the base of the $SL\nu$ consideration accounting for the plasma effects is the presence of the plasmon mass in the angular distribution. Note that the presence of plasmon mass makes the $SL\nu$ angular distribution analogous to the angular distribution of on-flight decay of a massive particle. As it follows from the exact expressions that govern the angular distribution on Fig.1 and Fig.2, the $SL\nu$ angular distribution goes to infinity as the angle reaches its boundary value (20) that corresponds to the outer cone on Fig.1. However the distribution is integrable and the contribution of the divergent areas into the total radiation power is negligible as it is common for kinematics of nuclear reactions (see [16]).

4. Conclusions

We developed a detailed evaluation of the spin light of neutrino in matter accounting for effects of the emitted plasmon mass. On the base of the exact solution of the modified Dirac equation for the neutrino wave function in the presence of the background matter the appearance of the threshold for the considered process is confirmed. The obtained exact and explicit threshold condition relation exhibit a rather complicated dependance on the matter density and neutrino mass. The dependance of the rate and power on the neutrino energy, matter density and the angular distribution of the $SL\nu$ is investigated in details. It is shown how the rate and power wash out when the threshold parameter $a = m_\gamma^2/4\widetilde{n}p$ approaching unity. From the performed detailed analysis it is shown that the $SL\nu$ mechanism is practically insensitive to the emitted plasmon mass for very high densities of matter (even up to $n = 10^{41} cm^{-3}$) for ultra-high energy neutrinos for a wide range of energies starting from $E = 1$ TeV. This conclusion is of interest for astrophysical applications of $SL\nu$ radiation mechanism in light of the recently reported hints of $1 \div 10$

PeV neutrinos observed by IceCube [17].

5. Acknowledgments

This study has been partially supported by the Russian Foundation for Basic Research (grants N. 11-02-01509 and N. 12-02-06833) and the Ministry of Education and Science of Russia (state contract N. 12.741.11.0180 and projects N. 2012-1.2.1-12-000-1012-1958 and N. 2012-1.1-12-000-1011-6097).

References

[1] C. Giunti, A. Studenikin, *Phys.Atom.Nucl.* **72** (2009) 2151 , C. Broggini, C. Giunti, A. Studenikin, arXiv:1207.3980 [to appear in Advances in High Energy Physics].

[2] A. Lobanov and A. Studenikin, *Phys.Lett.B* **564** (2003) 27 , A. Lobanov and A. Studenikin, *Phys.Lett.B* **601** (2004) 171.

[3] A. Studenikin, A. Ternov, *Phys.Lett.B* **608** (2005) 107 , A. Grigorev, A. Studenikin, A. Ternov, *Phys.Lett.B* **622** (2005) 199.

[4] A. Lobanov *Phys.Lett.B* **619** (2005) 136.

[5] A. Studenikin, *J. Phys. A: Math. Gen.* **39** (2006) 6769; A. Studenikin, *J. Phys. A: Math. Theor.* **41** (2008) 164047.

[6] I. Balantsev, Yu. Popov, A. Studenikin, *J.Phys.A: Math.Theor.* **44** (2011) 255301.

[7] A.A. Sokolov, I.M. Ternov, *Sov.Phys.Dokl.* **8** (1964) 1203, *Dokl.Akad.Nauk Ser.Fiz.* **153** (1964) 1052.

[8] A.V. Kuznetsov, N.V. Mikheev, *Int.J.Mod.Phys.A* **22** (2007) 3211.

[9] R. Belvedere, D. Pugliese, J.A. Rueda, R. Ruffini, S.-S. Xue, *Nucl.Phys.A* **883** (2012) 1.

[10] A. Grigoriev, A. Lokhov, A. Studenikin, A. Ternov, in: Quantum field theory under the influence of external conditions , ed. by K.A. Milton, M. Bordag, World Scientific, 2010, p. 528 (proceedings of the QFEXT09, September, 2009, University of Oklahoma, USA), arXiv:1003.0630.

[11] A.V. Grigoriev, A.V. Lokhov, A.I. Studenikin, A.I. Ternov, Abstracts of the 25th International Conference on Neutrino Physics and Astrophysics (June 2012, Kyoto, Japan) p.75.

[12] V.N. Asseev at al., *Phys.Rev.D* **84** (2011) 112003.

[13] E.W. Otten, C. Weinheimer, *Rept.Prog.Phys.* **71** (2008) 086201.

[14] R.L. Bowers, J.A. Campbell, R.L. Zimmerman, *Phys.Rev.D* **7** (1973) 2278, 2289.

[15] A.V. Kuznetsov, N.V. Mikheev, A.M. Shitova, *Int. J. Mod. Phys.A* **26** (2011) 4773.

[16] A.M. Baldin, V.I. Goldaskii, I.L. Rozenthal, Kinematics of Nuclear Reactions, Oxford-London-N.Y.-Paris, Pergamon Press, 1961.

[17] A. Ishihara (on behalf of IceCUBE collaboration), presentation at the 25th International Conference on Neutrino Physics and Astrophysics (June 2012, Kyoto, Japan).